

Obstructions for generalized graphmanifolds to be nonpositively curved

P. Svetlov*

December 13, 2004

Abstract

An n -dimensional manifold M ($n \geq 3$) is called *generalized graph manifold* if it is glued of blocks that are trivial bundles of $(n-2)$ -tori over compact surfaces (of negative Euler characteristic) with boundary. In this paper two obstructions for generalized graph manifold to be nonpositively curved are described.

Each 3-dimensional generalized graph manifold with boundary carries a metric of nonpositive sectional curvature in which the boundary is flat and geodesic (B. Leeb). The last part of this paper contains an example of 4-dimensional generalized graph manifold with boundary, which does not admit a metric of nonpositive sectional curvature with flat and geodesic boundary.

0 Introduction

0.1 Generalized graph manifolds. Let $n \geq 3$. A generalized graph manifold is a compact n -dimensional manifold consisting of finitely many blocks

$$M = \bigcup_{v \in V} M_v.$$

We require that the following conditions 1-3 to be fulfilled.

*Partially supported by RFBR (grants nos. 02-01-00090) and by the grant NSH-1914.2003.1

1. Each block M_v , $v \in V$ is homeomorphic to $T^{n-2} \times S_v$, where T^{n-2} is the $(n-2)$ -dimensional torus and S_v is a compact oriented surface with boundary and with negative Euler characteristic.
2. The manifold M is glued from its blocks by diffeomorphisms between their boundary $(n-1)$ -tori.
3. The gluing diffeomorphisms do not identify the homotopy classes of the fiber $(n-2)$ -tori.

Note that the decomposition is unique (up to isotopy). If $n = 3$ our definition gives a class $\mathfrak{M}_0$ of irreducible Haken graph manifolds. Each irreducible Haken graph manifold M has a covering manifold $\widetilde{M} \in \mathfrak{M}_0$ (see [1]).

0.2 Geometrization of a block. Let $M_v \simeq T^{n-2} \times S_v$ be a block in a n -dimensional generalized graph manifold M . A metric of nonpositive sectional curvature on M_v is said to be *geometrization* of M_v if

1. all boundary $(n-1)$ -tori of M_v are flat and geodesic;
2. all fiber $(n-2)$ -tori in M_v are flat, geodesic and isometric.

A block is geometrizable if and only if it admits a metric of nonpositive sectional curvature in which its boundary tori are flat and geodesic (see [2], and [3]).

0.3 Geometrization of a manifold. We say that a generalized graph manifold M is *geometrizable* if it admits a metric of nonpositive sectional curvature which induced a geometrization on each its block. A generalized graph manifold M is geometrizable if and only if it admits a metric of nonpositive sectional curvature in which its boundary tori are flat and geodesic (if M is closed, then it is geometrizable iff it admits a metric of nonpositive sectional curvature, see [2], and [3]).

In [2], S. V. Buyalo and V. L. Kobel'skii found necessary and sufficient conditions for closed generalized graph manifolds to carry a metric of nonpositive sectional curvature.

0.4 Results. In this paper two topological invariants $c(M), c'(M) \in \mathbb{Z}$ of generalized graph manifold M are described (see 3.1 and 4.1). If any of the numbers is nonpositive then M does not admit a geometrization.

Each 3-dimensional generalized graph manifold with boundary admits a geometrization (B. Leeb). The last section contains an example of 4-dimensional generalized graph manifold with boundary, which does not admit a geometrization (theorem 5.1).

Acknowledgements. The author is grateful to professor S. Buyalo for useful discussions.

1 Some notations

Let M be a generalized graph manifold and let V be the set of its blocks. Let $v \in V$. We define a set ∂v as the set of boundary components of the block M_v :

$$\partial M_v = \bigcup_{w \in \partial v} T_w$$

(each T_w is $(n-1)$ -dimensional torus). Now we divide all boundary tori in two classes:

$$W = \left\{ w \in \bigcup_{v \in V} \partial v \mid T_w \not\subset \partial M \right\}, \quad W_\partial = \left\{ w \in \bigcup_{v \in V} \partial v \mid T_w \subset \partial M \right\}.$$

A torus T_w , $w \in W$ is called *gluing torus*. The gluing diffeomorphisms induce a permutation $\varphi : W \rightarrow W$ (the torus T_w , $w \in W$ is glued with torus $T_{\varphi(w)}$ in M). If $w \in W$, we use the notation $-w$ for $\varphi(w)$ and $\psi_w : T_w \rightarrow T_{-w}$ for the gluing diffeomorphism.

2 Geometrization of a block

2.1 Waldhausen bases. Let $L_w = H_1(T_w, \mathbb{Z}) \simeq \mathbb{Z}^{n-1}$, $w \in \partial v$ and $\iota_v^w : F_v \rightarrow L_w$ be the inclusion of the first homology group of a fiber of the block $M_v = T^{n-2} \times S_v$. Note, that $\iota_v^w(F_v) = F_w$ is a maximal subgroup of L_w . A set $\{z_w \in L_w \mid w \in \partial v\}$ represents a boundary if

$$\sum_{w \in \partial v} (\iota_w)_* z_w = 0 \in H_1(M_v, \mathbb{Z}),$$

where $\iota_w : T_w \rightarrow M_v$ is the inclusion $T_w \subset \partial M_v \subset M_v$.

Let $(f_v^1, \dots, f_v^{n-2})$ be a basis in F_v and let a set $\{z_w \in L_w \mid w \in \partial v\}$ represents a boundary. Then for each $w \in \partial v$ the set

$$(z_w, f_w) = (z_w, \iota_v^w f_v^1, \dots, \iota_v^w f_v^{n-2})$$

is a basis in L_w . The set $\{(z_w, f_w) \mid w \in \partial v\}$ is called Waldhausen basis of the block M_v . If $(\bar{f}_v^1, \dots, \bar{f}_v^{n-2})$ is another basis in F_v and a set $\{\bar{z}_w \in L_w \mid w \in \partial v\}$ represents a boundary, then there exists a set of matrices

$$h_w = \begin{pmatrix} \det \sigma_v & 0 \\ n_w & \sigma_v \end{pmatrix} \in SL_{n-1}(\mathbb{Z}), \quad w \in \partial v$$

such that

$$(\bar{z}_w, \bar{f}_w) = (z_w, f_w) \cdot h_w = (\det \sigma_v \cdot z_w + f_w n_w, f_w \sigma_v)$$

for each $w \in \partial v$ (here $\bar{f}_w = (\iota_v^w \bar{f}_v^1, \dots, \iota_v^w \bar{f}_v^{n-2})$). It is clear that $\sum_{w \in \partial v} n_w = 0$ (see [2]).

2.2 The Leeb conditions. Each geometrization of M_v defines (and is defined by) a set of positively definite quadratic forms $\{\beta_w : L_w \rightarrow \mathbb{R} \mid w \in \partial v\}$, such that

L1) the map $\beta_w \circ \iota_v^w : F_v \rightarrow \mathbb{R}$ does not depend on $w \in \partial v$;

L2) if a set of classes $\{z_w \in L_w \mid w \in \partial v\}$ represents a boundary then

$$\sum_{w \in \partial v} B_w(z_w, \iota_v^w h) = 0 \quad \text{for any } h \in F_v$$

(here B_w is the bilinear form associated with β_w). The conditions L1,2 are called *the Leeb conditions* (see [2], and [3]). We note that the condition L2 is correct: if it is true for some set $\{z_w \in L_w\}$ represented a boundary then so is for each set $\{\bar{z}_w \in L_w\}$ represented a boundary:

$$\begin{aligned} \sum_{w \in \partial v} B_w(\bar{z}_w, \iota_v^w h) &= \sum_{w \in \partial v} B_w(\det \sigma_v \cdot z_w + f_w n_w, \iota_v^w h) = \\ &= (\det \sigma_v) \sum_{w \in \partial v} B_w(z_w, \iota_v^w h) + B_v(f_v, h) \sum_{w \in \partial v} n_w = 0. \end{aligned}$$

Here $B_v = (\iota_v^w)^* B_w$ (by L1 it is correctly defined).

2.3 The Gram matrix. Let $(z, f)_v = \{(z_w, f_w) \mid w \in \partial v\}$ be a Waldhausen basis of the block M_v . We define a set of *Gram matrices* $\{G_w^{(z, f)_v} \mid w \in \partial v\}$ for the geometrization $\{\beta_w\}_{w \in \partial v}$ in the basis $(z, f)_v$ as follows:

$$G_w^{(z, f)_v} = \begin{pmatrix} x_w & l_w \\ l_w^T & G_v \end{pmatrix},$$

where the number x_w , the $(n-2)$ -row l_w , and the square matrix G_v are defined by

$$x_w = B_w(z_w, z_w), \quad (l_w)_i = B_w(z_w, \iota_v^w f_v^i), \quad (G_v)_{ij} = B_v(f_v^i, f_v^j).$$

The condition L2 gives

$$\sum_{w \in \partial v} l_w = \sum_{w \in \partial v} B_w(z_w, l_v^w f_v) = 0,$$

so the row $\sum_w l_w$ is zero. It is clear, that if $(\bar{z}_w, \bar{f}_w) = (z_w, f_w) \cdot h_w$ is another Waldhausen basis, then $G_w^{(\bar{z}, \bar{f})_v} = h_w^T G_w^{(z, f)_v} h_w$.

3 The first obstruction

Let M be a n -dimensional generalized graph manifold, let $M_v, M_{v'}$ be blocks in M and let $\psi_w : T_w \rightarrow T_{-w}$ be the gluing map (here $w \in \partial v, -w \in \partial v'$). If the generalized graph manifold M has a geometrization (as it described in 2.2) then the induced maps $\Psi_w : (L_w, \beta_w) \rightarrow (L_{-w}, \beta_{-w})$

$$(z_{-w}, f_{-w}) = (z_w, f_w) \Psi_w$$

are isometries. Indeed, the homogeneous system of linear equations

$$\begin{cases} G_{-w} = \Psi_w^T G_w \Psi_w, & w \in W \\ \sum_{w \in \partial v} l_w = 0, & v \in V \end{cases} \quad LS_1(M)$$

has a nontrivial solution

$$G_w = \begin{pmatrix} x_w & l_w \\ l_w^T & G_v \end{pmatrix}, \quad w \in \partial v, \quad v \in V.$$

Let $c(M, (z, f))$ — be corank of the linear system (the number of its variables minus its rank). Note that a homogeneous linear system has a nontrivial solution iff its corank is positive.

Theorem 3.1. *The integer $c(M, (z, f))$ does not depend on the choice of the basis (z, f) .*

Proof. If we change Waldhausen bases in blocks of M then the matrix A of $LS_1(M)$ changes to a matrix AH , where H is nonsingular matrix. $\square$

So $c(M) = c(M, (z, f))$ is a topological invariant of M . Note, that if $c(M) \leq 0$ then M does not admit geometrization.

Lemma 3.2. *If M is a n -dimensional generalized graph manifold M then*

$$c(M) \geq (n-1)|W_\partial| + \frac{(n-2)(n-3)}{2} \cdot |V| - \frac{(n-1)(n-4)}{4} \cdot |W|.$$

Proof. By direct calculation with using the obvious inequality between the rank and the number of equations in the linear system. $\square$

If $n = 3, 4$ then $c(M) > 0$ for any n -dimensional generalized graph manifold M . So the obstruction to geometrization vanishes.

4 The second obstruction

Let M be a n -dimensional generalized graph manifold, let $M_v, M_{v'}$ be blocks in M and let $\psi_w : T_w \rightarrow T_{-w}$ be the gluing map (here $w \in \partial v, -w \in \partial v'$). Assume that M has a geometrization (as it described in 2.2). Then the induced maps $\Psi_w : (L_w, \beta_w) \rightarrow (L_{-w}, \beta_{-w})$

$$(z_{-w}, f_{-w}) = (z_w, f_w)\Psi_w$$

are isometries.

Let $n > 3$. Then the intersection

$$I_w = \Psi_{-w}(\iota_{v'}^{-w} F_{v'}) \cap \iota_v^w F_v$$

is non-empty $(n-3)$ -dimensional sublattice in L_w . The restriction

$$\Psi_w|_{I_w} : (I_w, \beta_w|_{I_w}) \rightarrow (I_{-w}, \beta_{-w}|_{I_{-w}})$$

must be isometry. Let

$$\Psi_w = \begin{pmatrix} a_w & b_w \\ c_w & d_w \end{pmatrix}$$

be the matrix in some Waldhausen bases, and let P_w be a nonzero $(n-2) \times (n-3)$ matrix of rank $n-3$ s.t. $b_w P_w = 0$. The equality $f_{-w} = z_w b_w + f_w d_w$ implies $f_{-w} P_w = f_w d_w P_w$. The vectors consist a basis of $I_w \simeq I_{-w}$, so we have

$$P_w^T (G_{v'} - d_w^T G_v d_w) P_w = 0 \quad LS_2(M)$$

Let $c'(M, (z, f))$ — be corank of the linear system (the number of its variables minus its rank). Note that a linear system has a nontrivial solution iff its corank is positive.

The following two proposition are proved in the same way as 3.1 and 3.2.

Theorem 4.1. *The integer $c'(M, (z, f))$ does not depend on the choice of the basis (z, f) .*

So $c'(M) = c'(M, (z, f))$ is a topological invariant of M . Note, that if $c'(M) \leq 0$ then M does not admit geometrization.

Lemma 4.2. *If M is a n -dimensional generalized graph manifold M and $n > 3$ then*

$$c'(M) \geq \frac{(n-2)(n-1)}{2} \cdot |V| - \frac{(n-3)(n-2)}{4} \cdot |W|.$$

5 The example

Let M be a generalized graph manifold with one boundary torus pasted from one block $M_1 = S \times T^2$ (here S is an oriented surface with 7 boundary circles) along gluing maps $\psi_i : T_i \rightarrow T_{-i}$, $i = 1, 2, 3$, where

$$\partial M_1 = \bigcup_{i=-3}^3 T_i, \quad \text{and} \quad \Psi_i = \begin{pmatrix} 1 & 1 & i-1 \\ 0 & 1 & 1 \\ 0 & 3 & 2 \end{pmatrix}, \quad i = 1, 2, 3$$

(in some Waldhausen basis). Note, that $c'(M) \geq 0$ (lemma 4.2)

Theorem 5.1. *The generalized graph manifold M with boundary $\partial M = T_0$ does not admit a geometrization.*

Proof. The linear system $LS_2(M)$ is

$$P_i^T (G_v - d_i^T G_v d_i) P_i = 0, \quad i = 1, 2, 3 \quad (*)$$

where

$$d_i = \begin{pmatrix} 1 & 1 \\ 3 & 2 \end{pmatrix}, \quad \text{and} \quad P_i = \begin{pmatrix} i-2 \\ 3i-1 \end{pmatrix}.$$

So, the system $(*)$ can be presented as follows:

$$(15i^2 - 52i + 45)x + (66i^2 - 232i + 204)y + (72i^2 - 258i + 231)z = 0, \quad i = 1, 2, 3.$$

(here x, y, z are entries of G_v). The matrix of $(*)$ is

$$A = \begin{pmatrix} 8 & 38 & 45 \\ 1 & 4 & 3 \\ 24 & 102 & 105 \end{pmatrix}$$

and rank of A is 3, so $c'(M) = 0$. Therefore the manifold M does not admit a geometrization. $\square$

- [1] S. Buyalo, V. Kobel'skii, *Geometrization of graphmanifolds. I. Conformal geometrization*, St. Petersburg Math. J., **7** (1996), No. 2, 185-216
- [2] S. Buyalo, V. Kobel'skii, *Generalized graphmanifolds of nonpositive curvature*, St. Petersburg Math. J., **11** (2000), No. 2, 251-268
- [3] B. Leeb, *3-manifolds with(out) metrics of nonpositive curvature*, Invent. Math., **122**, (1995), 277-289

St. Petersburg Branch,
Steklov Mathematical Institute,
Fontanka 27, 191023 St.Petersburg, Russia

svetlov@pdmi.ras.ru